

10.0 BLOOD DRUGS AND BLOOD-FORMING AGENTS

10.1 ANTI-ANEMICS

FERROUS SALT

Oral: tablet, equivalent to 60 mg
elemental iron solution, 15 mg
elemental iron/0.6 mL drops,
15 mL and 30 mL
30 mg elemental iron/5 mL
syrup, 60 mL

N.B.: The elemental iron content of
a ferrous salt depends on the
type of preparation as follows:

Ferrous fumarate	- 33%
Ferrous gluconate	- 12%
Ferrous lactate	- 19%
Ferrous sulfate, hydrated	- 20%
Ferrous sulfate, desiccated	- 32%

FOLIC ACID

Oral: 1 mg tablet
5 mg tablet

★ Inj.: 1 mg/mL, 1 mL ampul (IM)
(as sodium salt)

★ *HYDROXOCOBALAMIN*
(vitamin B12)

Oral: 100 mcg and 250 mcg tablet
Inj.: 1 mg/mL, 1 mL ampul/vial (IM)

mecobalamin

Oral: 500 mcg tablet
Inj.: 500 mcg/mL ampul (IM, IV)

10.2 HEMATOPOIETIC GROWTH FACTORS

epoetin alfa
(recombinant
human
erythropoietin)
(1, 2)

Inj.: 2,000 IU/0.2 mL, pre-filled syringe (IV)
2,000 IU/0.5 mL, pre-filled syringe (IV)
4,000 IU/0.4 mL, pre-filled syringe (IV)
10,000 IU/mL, pre-filled syringe (IV)
1,000 IU/0.5 mL, pre-filled syringe
(HSA-free) (IV)
2,000 IU/0.5 mL, pre-filled syringe
(HSA-free) (IV)
▼ 3,000 IU/0.3 mL, pre-filled syringe
(HSA-free) (IV)
4,000 IU/0.4 mL, pre-filled syringe
(HSA-free) (IV)
10,000 IU/mL, pre-filled syringe
(HSA-free) (IV)

epoetin beta
(recombinant
erythropoietin)
(1, 2)

Inj.: 2,000 IU/0.3 mL, pre-filled syringe
with needle (IV, SC)
4,000 IU/0.3 mL, pre-filled syringe
with needle (IV, SC)
5,000 IU/0.3 mL, pre-filled syringe
with needle (IV, SC)
lyophilized powder, 1,000 IU vial +
solvent ampul (IV, SC)
lyophilized powder, 10,000 IU vial +
solvent ampul (IV, SC)

filgrastim
(G-CSF)
(1, 2)

Inj.: 150 mcg/0.6 mL, vial (IV, SC)
300 mcg/mL, vial (IV, SC)
300 mcg/1.2 mL, vial (IV, SC)

★ *molgramostim*
(Gm-CSF)
(1, 2)

Inj.: 150 mcg and 400 mcg vial (IV, SC)

10.3 ANTICOAGULANTS

HEPARIN

LOW MOLECULAR WEIGHT HEPARIN

dalteparin (1, 2)

Inj.: 2,500 IU/0.2 mL and 5,000 IU/0.2 mL,
pre-filled syringe (SC)
(as sodium)
10,000 IU/1 mL, 1 mL ampul (SC)
(as sodium)

▼ *enoxaparin* (1, 2)

Inj.: 20 mg/0.2 mL, 40 mg/0.4 mL,
60 mg/0.6 mL and 80 mg/0.8 mL
pre-filled syringe (SC)
(as sodium)
100 mg/mL, 1 mL vial (SC)
(as sodium)

nadroparin (1, 2)

Inj.: 950 IU/0.1 mL, 0.2 mL, 0.3 mL,
0.4 mL, 0.6 mL, 0.8 mL, and
1 mL pre-filled syringe (SC)
(as calcium)

UNFRACTIONATED HEPARIN (1, 2)

Inj.: 1,000 IU/mL and 5,000 IU/mL, 5 mL
vial (IV, IV Infusion, SC)
(as sodium salt)

WARFARIN (1, 2)

Oral: 1 mg, 2.5 mg and 5 mg tablet
(as sodium salt)

10.4 ANTITHROMBOTICS (ANTIPLATELETS)

ASPIRIN

Oral: 80 mg and 325 mg tablet

▼ *clopidogrel*

Oral: 75 mg tablet

dipyridamole (2)Oral: 25 mg, 50 mg and 75 mg tablet
(preferably used in
combination with aspirin)

10.5 THROMBOLYTIC (FIBRINOLYTIC)

STREPTOKINASE (1)

Inj.: powder, 750,000 IU vial (IV Infusion)
and 1,500,000 IU vial
(IV Infusion)★ *recombinant tissue
plasminogen
activator (rTPA)*
(1, 2)

Inj.: 1 mg/mL 50 mL vial (IV Infusion)

10.6 ANTI-FIBRINOLYTIC

*tranexamic acid*Oral: 250 mg and 500 mg capsule
500 mg tabletInj.: 100 mg/mL, 2.5 mL and 5 mL
ampul (IM, IV)