

## 11.0 BLOOD PRODUCTS AND BLOOD SUBSTITUTES

### 11.1 PLASMA EXPANDERS / SUBSTITUTES (COLLOIDS)

<i>DEXTRAN, LOW MOLECULAR WEIGHT (dextran 40)</i>	Inj.:	10% dextran 40 in 0.9% sodium chloride, 500 mL bottle (IV Infusion) 10% dextran 40 in 5% dextrose, 500 mL bottle (IV Infusion)
<i>dextran, high molecular weight (dextran 70)</i>	Inj.:	6% dextran 70 in 0.9% sodium chloride, 500 mL bottle (IV Infusion) 6% dextran 70 in 5% dextrose, 500 mL bottle (IV Infusion)
<i>hydroxyethyl starch</i>	Inj.:	6% solution, 250 mL and 500 mL bottle (IV Infusion) 10% solution, 250 mL and 500 mL bottle (IV Infusion)
<i>modified fluid gelatin (polymerisate of degraded succinylated gelatin)</i>	Inj.:	3% and 4% solution, 500 mL bottle (IV Infusion)
★ <i>polygeline</i>	Inj.:	3.5% colloidal solution, 500 mL bottle (IV Infusion)

### 11.2 PLASMA FRACTIONS FOR SPECIFIC USES

(All plasma fractions should comply with the WHO requirements for the Collection, Processing and Quality Control of Human Blood and Blood Products)

<i>ALBUMIN, HUMAN (1, 2)</i>	Inj.:	20%, 50 mL and 100 mL bottle (IV, IV Infusion) 25%, 50 mL and 100 mL bottle (IV, IV Infusion)
<i>FACTOR VIII CONCENTRATE (1, 2)</i>	Inj.:	lyophilized powder, 100 IU/g vial + diluent (IV)
<i>FACTOR IX COMPLEX CONCENTRATE (coagulation factors II, VII, IX, X) (1, 2)</i>	Inj.:	100 IU/mL, 5 mL and 10 mL vial (IV) lyophilized powder, 500 IU vial + diluent (IV)